

JAN 13 1975

UNIVERSITY OF CALIFORNIA, BERKELEY

BERKELEY • DAVIS • IRVINE • LOS ANGELES • RIVERSIDE • SAN DIEGO • SAN FRANCISCO



SANTA BARBARA • SANTA CRUZ

VIRUS LABORATORY

Department of Molecular Biology

BERKELEY, CALIFORNIA 94720

January 8, 1975

Dr. Joshua Lederberg
Department of Genetics
Stanford University School of Medicine
Stanford, California 94305

Dear Josh:

Cal Ward must be referring to a Special Topics Course on RNA and DNA regulation in bacteria I taught in the Spring of 1969, as my farewell to Molecular Genetics just before taking my Neurobiological orders at Harvard Medical School. I am afraid my memory is very hazy and I am not sure I have my notes any longer. I am sure I didn't give any empirical data on maximal growth rates.

What I think I did was to calculate the maximum rate of r-RNA synthesis per DNA genome, assuming that all the r-RNA cistrons are densely packed with RNA polymerase molecules, and using the RNA chain growth step time we had then recently measured in vivo. If this maximum rate is q , then at a rate of DNA synthesis r per genome, the maximum ratio of RNA/DNA is q/r . Now using Maaløe's empirical linear relation between [RNA/DNA] content of bacteria and r (where r is inversely related to the generation time via the genome size) you can find the value of r where q/r lies on the Maaløe line. That value of r gives you some kind of theoretical minimum generation time (if one assumes, as is nearly true, that the RNA chain growth rate is invariant).

I am sorry that this is the best I can do "at this point in time". I would suggest that you contact Ole in this matter, who had been my guiding beacon in the first place. Meanwhile, I send you my warmest regards

Your

Gunther S. Stent

GSS:mh